arXiv:1601.03515v1 [math.GN] 14 Jan 2016

ON $I_{\alpha^m}(A)$, $C_{\alpha^m}(A)$ AND α^m-OPEN AND CLOSED FUNCTIONS

R. PARIMELAZHAGAN AND MILBY MATHEW

Abstract. In this paper, we derive more on α^m-continuous functions and α^m-irresolute functions with α^m-open maps and α^m-closed maps in topological spaces also we introduce $I_{\alpha^m}(A)$ and $C_{\alpha^m}(A)$ by using the α^m-open sets and α^m-closed sets and studied some of their properties.

Keywords : $I_{\alpha^m}(A)$ and $C_{\alpha^m}(A)$.

2000 Mathematics Subject Classification : 57N40.

1. Introduction

Malghan [9] introduced the concept of generalized closed maps in topological spaces. Devi [3] introduced and studied sg-closed maps and gs-closed maps. wg-closed maps and rwg-closed maps were introduced and studied by Nagavani [10]. Regular closed maps, gpr-closed maps and rg-closed maps have been introduced and studied by Long [7], Gnanambal [4] and Arockiarani [2] respectively.

In 2014, [12] we introduced the concepts of α^m-closed sets and α^m-open set in topological spaces. Also we have introduced the concepts of α^m-continuous functiopns and α^m-irresolute functions. In this paper, by deriving the properties of α^m-continuous functions and α^m-irresolute functions with α^m-open maps and α^m-closed maps in topological spaces. Further we introduce the concept of $I_{\alpha^m}(A)$ and $C_{\alpha^m}(A)$ by using the α^m-open sets and α^m-closed sets respectively. Further various characterisation are studied. are introduced

2. Preliminaries

Throughout this paper (X, τ) and (Y, σ) represent topological spaces on which no separation axioms are assumed unless otherwise mentioned. For a subset A of a space (X, τ), $cl(A)$, $int(A)$ and A^c denote the closure of A, the interior of A and the complement of A in X, respectively.

Definition 2.1. *A subset A of a topological space (X, τ) is called*

(a) *a preopen set* [9] *if $A \subseteq int(cl(A))$ and pre-closed set if $cl(int(A)) \subseteq A$.*

(b) *a semiopen set* [5] *if $A \subseteq cl(int(A))$ and semi closed set if $int(cl(A)) \subseteq A$.*

(c) *an α-open set* [11] *if $A \subseteq int(cl(int(A)))$ and an α-closed set if $cl(int(cl(A))) \subseteq A$.*

(d) a semi-preopen set [1] *(β-open set) if* $A \subseteq cl(int(cl(A)))$ *and semi-preclosed set if* $int(cl(int(A))) \subseteq A$.

(e) an α^m*-closed set* [12] *if* $int(cl(A)) \subseteq U$, *whenever* $A \subseteq U$ *and* U *is* α*-open. The complement of* α^m*-closed set is called an* α^m*-open set.*

Definition 2.2. *A space* (X, τ_X) *is called a* $T_{\frac{1}{2}}$*-space* [6] *if every g-closed set is closed.*

Definition 2.3. *A function* $f : (X, \tau) \longrightarrow (Y, \sigma)$ *is called*

(a) an α^m*-continuous* [13] *if* $f^{-1}(V)$ *is* α^m*-closed in* (X, τ) *for every closed set* V *of* (Y, σ).

(b) an α^m*-irresolute* [13] *if* $f^{-1}(V)$ *is* α^m*-closed in* (X, τ) *for every* α^m*-closed set* V *of* (Y, σ).
(b) an α*-irresolute* [11] *if* $f^{-1}(V)$ *is* α*-closed in* (X, τ) *for every* α*-closed set* V *of* (Y, σ).

3. More on α^m-open sets

Theorem 3.1. *A map* $f : (X, \tau) \longrightarrow (Y, \sigma)$ *is* α^m*-closed if and only if for each subset* S *of* (Y, σ) *and each open set* U *containing* $f^{-1}(S)$ *there is an* α^m*-open set* V *of* (Y, σ) *such that* $S \subseteq V$ *and* $f^{-1}(V) \subseteq U$.

Proof. Suppose f is α^m-closed. Let $S \subseteq Y$ and U be an open set of (X, τ) such that $f^{-1}(S) \subseteq U$. Then $V = (f(U^c))^c$ is an α^m-open set containing S such that $f^{-1}(V) \subseteq U$.

For the converse, let F be a closed set of (X, τ). Then $f^{-1}(f(F^c)) \subseteq F^c$ and F^c is open. By assumption, there exists an α^m-open set V in (Y, σ) such that $(f(F^c)) \subseteq V$ and $f^{-1}(V) \subseteq F^c$ and so $F \subseteq (f^{-1}(V))^c$. Hence $V^c \subseteq f(F) \subseteq f((f^{-1}(V))^c) \subseteq V^c$ which implies $f(F) = V^c$. Since V^c is α^m-closed, $f(F)$ is α^m-closed and therefore f is α^m-closed.

□

Proposition 3.2. *If* $f : (X, \tau) \longrightarrow (Y, \sigma)$ *is* α*-irresolute* α^m*-closed and* A *is an* α^m*-closed subset of* (X, τ), *then* $f(A)$ *is* α^m*-closed in* (Y, σ).

Proof. Let U be an α-open set in (Y, σ) such that $f(A) \subseteq U$. Since f is α-irresolute, $f^{-1}(U)$ is an α-open set containing A. Hence $cl(A) \subseteq f^{-1}(U)$ as A is α^m-closed in (X, τ). Since f is α^m-closed, $f(cl(A))$ is an α^m-closed set contained in the α-open set U, which implies that $cl(f(cl(A))) \subseteq U$ and hence $cl(f(A)) \subseteq U$. Therefore, $f(A)$ is

an α^m-closed set in (Y, σ).

□

Corollary 3.3. *3.8. Let $f : (X, \tau) \longrightarrow (Y, \sigma)$ be an α^m-closed and $g : (Y, \sigma) \longrightarrow (Z, \eta)$ be α^m-closed and α-irresolute, then their composition $g \circ f : (X, \tau) \longrightarrow (Z, \eta)$ is α^m-closed.*

Proof. Let A be a closed set of (X, τ). Then by hypothesis $f(A)$ is an α^m-closed set in (Y, σ). Since g is both α^m-closed and α-irresolute by the above Proposition, $g(f(A)) = (g \circ f)(A)$ is α^m-closed in (Z, η) and therefore $g \circ f$ is α^m-closed.

□

Proposition 3.4. *Let $f : (X, \tau) \longrightarrow (Y, \sigma)$ be a closed map and $g : (Y, \sigma) \longrightarrow (Z, \eta)$ be an α^m-closed map, then their composition $g \circ f : (X, \tau) \longrightarrow (Z, \eta)$.*

Proof. Let A be a closed set of (X, τ). Then by assumption $f(A)$ is closed in (Y, σ) and again by assumption $g(f(A))$ is α^m-closed in (Z, η). i.e., $(g \circ f)(A)$ is α^m-closed in (Z, η) and so $g \circ f$ is α^m-closed.

□

Theorem 3.5. *Let $f : (X, \tau) \longrightarrow (Y, \sigma)$ and $g : (Y, \sigma) \longrightarrow (Z, \eta)$ be two maps such that their composition $g \circ f : (X, \tau) \longrightarrow (Z, \eta)$ is an α^m-closed map. Then the following statements are true.*

(a) If f is continuous and surjective, then g is α^m-closed.

(b) If g is α^m-irresolute and injective, then f is α^m-closed.

Proof. (a) Let A be a closed set of (Y, σ). Since f is continuous, $f^{-1}(A)$ is closed in (X, τ) and since $g \circ f$ is α^m-closed, $(g \circ f)(f^{-1}(A))$ is α^m-closed in (Z, η). That is $g(A)$ is α^m-closed in (Z, η), since f is surjective. Therefore g is an α^m-closed map.

(b) Let B be a closed set of (X, τ). Since $g \circ f$ is α^m-closed, $(g \circ f)(B)$ is α^m-closed in (Z, η). Since g is α^m-irresolute, $g^{-1}((g \circ f)(B))$ is α^m-closed set in (Y, σ). That is $f(B)$ is α^m-closed in (Y, σ), since g is injective. Thus f is an α^m-closed map.

□

Definition 3.6. *(a) Let X be a topological space and let $x \in X$. A subset N of X is said to be α^m-nbbd of x if there exists an α^m-open set G such that $x \in G \subset N$.*

The collection of all α^m-nbhd of $x \in X$ is called an α^m-nbhd system at x and shall be denoted by $n^{\#_{\alpha^m}}(x)$.

(b) Let X be a topological space and A be a subset of X, A subset N of X is said to be α^m-nbhd of A if there exists an α^m-open set G such that $A \in G \subseteq N$.

(c) Let A be a subset of X. A point $x \in A$ is said to be an α^m-interior point of A, if A is an $n^{\#_{\alpha^m}}(x)$. The set of all α^m-interior points of A is called an α^m-interior of A and is denoted by $I_{\alpha^m}(A)$.

$I_{\alpha^m}(A) = \bigcup\{G : G \text{ is } \alpha^m\text{-open}, \ G \subset A\}$.

(d) *Let A be a subset of X. A point $x \in A$ is said to be an α^m-closure of A. Then $C_{\alpha^m}(A) = \bigcap\{F : A \subset F \in \alpha^m C(X, \tau_X)\}$.*

Theorem 3.7. *Let X be a topological space. If A is α^m-closed subset of X and $x \in C_{\alpha^m}(A)$ if and only if for any α^m-nbhd N of x in X, $N \cap A \neq \phi$.*

Proof. Let us assume that there is an α^m-nbhd N of the point x in X such that $N \cap A = \phi$. There exist an α^m-open set G of X such that $x \in G \subseteq N$. Therefore we have $G \cap A = \phi$ and so $x \in X - G$. Then $C_{\alpha^m}(A) \in X - G$ and therefore $x \notin C_{\alpha^m}(A)$, which is the contradiction to the hypothesis $x \in C_{\alpha^m}(A)$. Therefore $N \cap A \neq \phi$.

Conversely, Suppose that $x \notin C_{\alpha^m}(A)$, then there exists a α^m-closed set G of X such that $A \subseteq G$ and $x \notin G$. Thus $x \in X - G$ and $X - G$ is α^m-open in X and hence $X - G$ is an α^m-nbhd of x in X. But $A \cap (X - G) = \phi$ which is a contradiction. Hence $x \in C_{\alpha^m}(A)$.

□

Proposition 3.8. *If A be a subset of X, then $I_{\alpha^m}(A) = \bigcup\{G : G \text{ is } \alpha^m\text{-open}, \ G \subset A\}$.*

Proof. Let A be a subset of X, $x \in I_{\alpha^m}(A) \iff x$ is an α^m-interior point of A, A is an $n^{\#_{\alpha^m}}(x)$ which implies that there exists an α^m-open set G such that $x \in G \subset A$, $x \in \bigcup\{G : G \text{ is } \alpha^m\text{-open}, \ G \subset A\}$. Hence $I_{\alpha^m}(A) = \bigcup\{G : G \text{ is } \alpha^m\text{-open}, \ G \subset A\}$.

□

Theorem 3.9. *Assume that the collection of all α^m-open sets of Y is closed under arbitrary union. Let $f : (X, \tau) \longrightarrow (Y, \sigma)$ be a map. Then the following statements are equivalent:*

(a) *f is an α^m-open map.*

(b) *For a subset A of (X, τ), $f(int(A)) \subseteq I_{\alpha^m}(f(A))$.*

(c) *For each $x \in X$ and for each neighborhood U of x in (X, τ), there exists an α^m-nbhd W of $f(x)$ in (Y, σ) such that $W \subseteq f(U)$.*

Proof. (a) $\implies$ (b) Suppose f is α^m-open. Let $A \subseteq X$. Then $int(A)$ is open in (X, τ) and so $f(int(A))$ is α^m-open in (Y, σ). We have $f(int(A)) \subseteq f(A)$. Therefore, $f(int(A)) \subseteq I_{\alpha^m}(f(A))$.

(b) $\implies$ (c) Suppose (b) holds. Let $x \in X$ and U be an arbitrary neighborhood of x in (X, τ). Then there exists an open set G such that $x \in G \subseteq U$. By assumption, $f(G) = f(int(G)) \subseteq I_{\alpha^m}(f(G))$. This implies $f(G) = I_{\alpha^m}(f(G))$, we have

$f(G)$ is α^m-open in (Y,σ). Further, $f(x) \in f(G) \subseteq f(U)$ and so (c) holds, by taking $W = f(G)$.

$(c) \Longrightarrow (a)$ Suppose (c) holds. Let U be any open set in (X,τ)), $x \in U$ and $f(x) = y$. Then $y \in f(U)$ and for each $y \in f(U)$, by assumption there exists an α^m-nbhd W_y of y in (Y,σ) such that $W_y \subseteq f(U)$. Since W_y is an α^m-nbhd of y, there exists an α^m-open set V_y in (Y,σ) such that $y \in V_y \subseteq W_y$. Therefore, $f(U) = \cup\{V_y : y \in f(U)\}$ is an α^m-open set in (Y,σ) by the given condition. Thus f is an α^m-open map.

□

Corollary 3.10. *A map $f : (X,\tau) \longrightarrow (Y,\sigma)$ is α^m-open if and only if $f^{-1}(C_{\alpha^m}(B)) \subseteq cl(f^{-1}(B))$ for each subset B of (Y,σ).*

Proof. Suppose that f is α^m-open. Then for any $B \subseteq Y$, $f^{-1}(B) \subseteq cl(f^{-1}(B))$, there exists an α^m-closed set K of (Y,σ) such that $B \subseteq K$ and $f^{-1}(K) \subseteq cl(f^{-1}(B))$. Therefore, $f^{-1}(C_{\alpha^m}(B)) \subseteq (f^{-1}(K)) \subseteq cl(f^{-1}(B))$, since K is an α^m-closed set in (Y,σ).

Conversely, let S be any subset of (Y,σ) and F be any closed set containing $f^{-1}(S)$. Put $K = C_{\alpha^m}(S)$. Then K is an α^m-closed set and $S \subseteq K$. By assumption, $f^{-1}(K) = f^{-1}(C_{\alpha^m}(S)) \subseteq cl(f^{-1}(S)) \subseteq F$ and therefore, f is α^m-open.

□

References

[1] Andrijevic. D, Semi-preopen sets, Mat. Vesink, 38 (1986), 24 - 32.

[2] I. Arockiarani, Studies on Generalizations of Generalized Closed Sets and Maps in Topological Spaces, Ph. D Thesis, Bharathiar University, Coimbatore, (1997).

[3] R.Devi, Studies on generalizations of closed maps and homeomorphisms in topological spaces, Ph.D Thesis, Bharathiar University, Coimbatore, (1994).

[4] Y. Gnanambal, On Generalized Pre-regular Closed Sets in Topological Spaces, Indian J. Pure Appl. Math., 28(1997), 351-360.

[5] Levine. N, Semi-open sets and semi-continuity in topological spaces, Amer. Math. Monthly, 70 (1963), 36 - 41.

[6] Levine. N, Generalized closed sets in topology, Rend. Circ. Math. Palermo, 19 (1970), 89 - 96.

[7] P. E. Long and L. L. Herington, Basic Properties of Regular Closed Functions, Rend. Cir. Mat. Palermo, 27(1978), 20-28.

[8] S.R.Malghan, Generalized closed maps, J. Karnataka Univ. Sci., 27(1982), 82-88.

[9] Mashhour. A. S, Abd EI-Monsef. M. E. and EI-Deeb. S. N., On Precontinuous and weak pre-continuous mapping, Proc. Math., Phys. Soc. Egypt, 53 (1982), 47 - 53.

[10] N. Nagaveni, Studies on Generalizations of Homeomorphisms in Topological Spaces, Ph. D., Thesis, Bharathiar University, Coimbatore(1999).

[11] Njastad. O, On some classes of nearly open sets, Pacific J. Math., , 15 (1965), 961-970.

[12] Milby Mathew and R. Parimelazhagan, α^m-Closed Sets in Topological Spaces, International Journal of Mathematical Analysis Vol. 8, 2014, no. 47, 2325 - 2329.

[13] Milby Mathew and R. Parimelazhagan, α^m-Continuous Maps in Topological Spaces. (submitted)

Department of Mathematics, RVS Technical Campus, Coimbatore - 641 402, Tamilnadu, India. E-mail: pari_kce@yahoo.com.

Research Scholar, Department of Mathematics, Karpagam University, Coimbatore - 32, Tamilnadu, India. E-mail: milbymanish@yahoo.com